

KOI CARE
B - D - S
fluke
treatment


The skin and gill flukes *Dactylogyrus* Sp. and *Gyrodactylus* Sp. are notoriously difficult to treat directly in the pond. This is because they are protected by the natural mucus layer on the fish and by causing the fish to produce more mucus by irritating the skin and gills.

There is also some evidence that some species of these parasites are resistant to some regularly used treatments.

Successful treatment depends upon ingredients which can penetrate the mucus and reach the parasites.

B-D-S fluke treatment provides a well tried and tested formulation which treats both skin and gill flukes as well as bacterial diseases such as fin or tail-rot and ulcers in a strength sufficient to treat large ponds easily.

The correct dose of B-D-S fluke treatment does not affect the filter bacteria or plants and persists in the water long enough to destroy the various stages of parasite which may otherwise remain to re-infect treated fish.

- Remove adsorbent filter material - carbon, zeolite etc.
- Switch off UV
- Increase Aeration
- Allow plenty of time to observe fish
- Do not mix medications
- Leave 10 days between treatments if no improvement is seen

Dosage:

Add 10 ml of B-D-S to every 80 gallons (363 L) of pond water.

Pack size	Treats		
	Litres	Imp. Gallons	US Gallons
250 ml	9 000	2 000	2 380
500 ml	18 000	4 000	4 764
1000 ml	36 000	8 000	9 528
2500 ml	90 000	20 000	23 800